Lake County ENT/Head & Neck Specialists Phone (847) 662-4442 Fax (847) 662-4446

Post-operative Instructions following Thyroidectomy

General:

You may be hospitalized for one or two nights following your thyroid procedure. Your surgeon will see you each day of your hospitalization to ensure that you are recovering well. You should avoid non-steroidal anti-inflammatory drugs (NSAIDS) such as aspirin, ibuprofen, naproxen (Excedrin, Motrin, Naprosyn, Advil) for 10 days prior to the surgery. These drugs are mild blood thinners and will increase your chances of bleeding.

Diet:

You may have liquids by mouth once you have awakened from anesthesia. If you tolerate the liquids without significant nausea or vomiting then you may take solid foods without restrictions. Generally patients experience a mild sore throat for 2-3 days following thyroidectomy. This usually does not interfere with swallowing.

Pain Control:

Patients report mild to moderate neck pain for several days following thyroidectomy. This is usually well controlled with prescription strength oral pain medications such as Vicodin. Please take the pain medication prescribed by your surgeon when needed. You should avoid non-steroidal anti- inflammatory drugs (NSAIDS) such as aspirin, ibuprofen, naproxen (Excedrin, Motrin, Naprosyn, Advil) because these drugs are mild blood thinners and will increase your chances of having a post- operative bleed into the neck tissues or neck wound.

Activity:

Sleep with the head elevated for the first 48 hours. You may use two pillows to do this or sleep in a reclining chair. No heavy lifting or straining for 2 weeks following the surgery. You should plan for 1 week away from work. If your job requires manual labor, lifting or straining then you should be out of work for 2 weeks or limited to light duty until the 2 week mark.

Wound Care:

Do not wash or manipulate the neck wound for 48 hours following the thyroidectomy (except to apply ointment). The neck dressing will be removed on the morning following your surgery. If a drain has been placed in the neck, this will be removed prior to your discharge from the hospital unless otherwise directed by your surgeon. Once you are home, apply a thin layer of the Vaseline to the wound 3 times daily. You may shower and allow the wound to get wet 48 hours following the surgery. Allow soap and water to run over the wound. Do not scrub or manipulate the wound for 7 days. Pat the area dry, don't rub it with a towel. After 7 days you may gently lather the wound with soap and water.

(Continued \longrightarrow)

Follow-up Appointment:

Your follow-up appointment in the office will be 5-8 days following your surgery. At the post- operative visit, your sutures are removed and the pathology report is reviewed.

Please call our office immediately if you experience:

*Difficulty breathing *Neck swelling *Numbness or tingling in your fingertips or lips *Muscle cramps or spasms *Bleeding from the wound *Fever greater than 101 degrees Fahrenheit *Pus coming from the wound *Increasing redness around the wound

Office phone: (847) 662-4442